arXiv:1202.2051v1 [hep-ph] 9 Feb 2012

# Thermal pions in a magnetic background

Jens O. Andersen
*Department of Physics, Norwegian University of Science and Technology, N-7491 Trondheim, Norway and*
*Niels Bohr International Academy, Blegdamsvej 17,DK-2100 Copenhagen, Denmark*

(Dated: February 10, 2012)

We use chiral perturbation theory for $SU(2)$ to compute the leading corrections to the thermal mass of the pions and the pion decay constant in the presence of the magnetic field in a low-temperature expansion. The magnetic field gives rise to a splitting between $M_{\pi^0}$ and $M_{\pi^\pm}$ as well as $F_{\pi^0}$ and $F_{\pi^\pm}$. We also calculate the free energy and the quark condensate to next-to- leading order. The results suggest that the critical temperature $T_c$ for the chiral transition is larger in the presence of a constant magnetic field, in agreement with most model calculations but in disagreement with recent lattice calculations.

*Introduction*: Chiral perturbation theory (ChPT) provides a systematic framework for calculating properties of QCD at low energies [1–4]. ChPT is not an expansion in powers of some small coupling constant, but it is a systematic expansion in powers of momenta $p$ where a derivative counts as one power and the quark masses count as two powers. Chiral perturbation theory is a nonrenormalizable quantum field theory in the old sense of the word. This means that a calculation at a given order in momentum $p$, requires that one adds higher-order operators in order to cancel the divergences that arise in the calculations at that order. This implies that one needs more and more couplings and therefore more experiments to determine them. However, this poses no problem, as long as one is content with finite precision. This is is the essence of effective field theory. The chiral Lagrangian that describes the (pseudo)Goldstone bosons are uniquely determined by the global symmetries of QCD and the assumption of symmetry breaking. The Lagrangian $\mathcal{L}_{\rm eff}$ consists of a string of terms that involve an increasing number of derivatives or quark mass factors, each multiplied by a low-energy constant (LEC) $l_i$. However, QCD is a confining and strongly interacting theory at low energies. Thus the couplings $l_i$ of the chiral Lagrangian cannot be calculated from QCD. Instead, the couplings are fixed by experiments

The thermodynamics of a pion gas using ChPT was studied in detail in a series of papers 25 years ago [5–7]. The thermal pion mass and the thermal pion decay constant were evaluated at next-to-leading order, while the pressure and the temperature dependence of the quark condensate were calculated to next-to-next-to-leading order. In this letter, we present the calculation of these quantities in the presence of a constant magnetic background. The details of the calculations can be found in Ref. [8].

QCD in external magnetic fields has received a lot of attention in recent years due to its relevance in several physical situations. For example, large magnetic fields exist inside ordinary neutron stars as well as magnetars [9]. In the latter case, the cores may be color superconducting and so it is important to study the effects of external magnetic fields in this phase. Similarly, it has been suggested that strong magnetic fields are created in heavy-ion collisions at the Relativistic Heavy-Ion Collider (RHIC) and the Large Hadron Collider (LHC) and that these play an important role [10]. In this case, the magnetic field strength has been estimated to be up to $B \sim 10^{19}$ Gauss, which corresponds to $|qB| \sim 6m_\pi^2$, where $|q|$ is the charge of the pion. This has spurred the interest in studying QCD in external fields. At zero baryon chemical potential this can be done from first principles using lattice simulations and some recent result are found in [11–13].

Chiral perturbation theory has been used to study the quark condensate in strong magnetic fields at zero temperature [14–17], while the quark-hadron phase transition was investigated in Ref. [18]. The effects of external magnetic fields on the chiral transition have been studied in detail using the NJL model [19–28], the Polyakov-loop extended NJL model [29, 30], the quark-meson model [27, 28, 31, 32], the (P)QM model [33, 34], the linear sigma model [36] and the MIT bag model [35]

*Chiral perturbation theory:* As explained in the introduction, chiral perturbation theory is low-energy effective field theory that can be used to systematically calculate physical quantities as a power series in momentum. The effective Lagrangian is given by an infinite string of operators involving an increasing number of derivatives or quark masses. Schematically, we can write $\mathcal{L}_{\rm eff} = \mathcal{L}^{(2)} + \mathcal{L}^{(4)} + \mathcal{L}^{(6)} + ...$ where the superscript indicates the powers of momentum.

By expanding the Lagrangian $\mathcal{L}^{(2)}$ to fourth order in the pion fields $\pi_i$, we obtain

$$\begin{aligned}\mathcal{L}^{(2)} &= \frac{1}{2}\left(\partial_\mu\pi_0\right)^2+\frac{1}{2}M^2\pi_0^2+(\partial_\mu+iqA_\mu)\pi^+(\partial_\mu-iqA_\mu)\pi^-+M^2\pi^+\pi^--\frac{M^2}{24F^2}\left[(\pi^0)^2+2\pi^+\pi^-\right]^2\\&+\frac{1}{6F^2}\left[-2\pi_0^2(\partial_\mu\pi^+)(\partial_\mu\pi^-)-2\pi^+\pi^-(\partial_\mu\pi_0)^2+[\partial_\mu(\pi^+\pi^-)]^2+2\pi_0[\partial_\mu\pi_0][\partial_\mu(\pi^+\pi^-)]-4\pi^+\pi^-(\partial_\mu\pi^+)(\partial_\mu\pi^-)\right] ,\end{aligned}$$

where we have defined the complex pion fields as $\pi_\pm=\frac{1}{\sqrt{2}}(\pi_1\pm i\pi_2)$ and $A_\mu=B\delta_{\mu2}x_1$. Similarly, expanding $\mathcal{L}^{(4)}$ to second order in the pion fields yields

$$\begin{aligned}\mathcal{L}^{(4)} &= \frac{2l_5}{F^2}(qF_{\mu\nu})^2\pi^+\pi^--2il_6qF_{\mu\nu}\left[(\partial_\mu\pi^-)(\partial_\nu\pi^+)+iqA_\mu\partial_\nu(\pi^+\pi^-)\right]+(l_3+l_4)\frac{M^4}{F^2}\pi_0^2+2(l_3+l_4)\frac{M^4}{F^2}\pi^+\pi^-\\&+l_4\frac{M^2}{F^2}(\partial_\mu\pi_0)^2+2l_4\frac{M^2}{F^2}(D_\mu\pi^+)(D_\mu\pi^-) .\end{aligned}$$

The Lagrangian $\mathcal{L}^{(6)}$ is very complicated as it contains more than 50 terms for $SU(2)$ [4]. However, only one term is relevant for the present problem [15, 17], namely

$$\mathcal{L}^{(6),\text{relevant}}=4c_{34}M^2(qF_{\mu\nu})^2 .$$

We have used the parametrization $U=e^{i\pi_i\tau_i/F}$, where $\tau_i$ are the Pauli matrices. This is different from the parametrization used in [15, 16] and so the expressions for $\mathcal{L}$ also differ. However, we get the same result for physical quantities as we should.

The chiral Lagrangian comes with a number of undetermined parameters or low-energy constants. These parameters can be determined by experiments; however, loop corrections involve renormalization of them. The relation between the bare and renormalized parameters has been can be expressed as $l_i=-\frac{\gamma_i}{2(4\pi)^2}\left[\frac{1}{\epsilon}+1-\bar{l}_i\right]$, where $\gamma_i$ are coefficients and $\bar{l}_i$ are scale-independent parameters [2].

*Pion mass and pion decay constant:* The pion masses $M_{\pi^0}$ and $M_{\pi^\pm}$ are defined by the position of the pole of the propagator. At leading order, the expressions for them are divergent and requires renormalization of the parameters $l_3$, $l_5$, and $l_6$. The result is

$$\begin{aligned}M_{\pi^0}^2 &= M_\pi^2\left[1+\frac{1}{(4\pi)^2F^2}\left(I_H-\frac{1}{2}J_1T^2+J_1^B|qB|\right)\right] ,\\ M_{\pi^\pm} &= M_\pi^2\left[1+\frac{T^2}{2(4\pi)^2F^2}J_1\right]+\frac{(qB)^2}{3(4\pi)^2F^2}(\bar{l}_6-\bar{l}_5) ,\end{aligned}$$

where the pion mass $M_\pi^2$ in the vacuum is given by $M_\pi^2=M^2(1-\frac{M^4}{2(4\pi)^2F^2}\bar{l}_3)$, $I_H=M^2\log\frac{M^2}{2|qB|}-M^2-2\zeta^{(1,0)}(0,\frac{1}{2}+x)|qB|$, $J_1=4\beta^2\int_0^\infty p^2dp/\sqrt{p^2+M^2}(e^{\beta\sqrt{p^2+M^2}}-1)$, $J_1^B=2\sum_{m=0}^\infty\int_0^\infty dp/\sqrt{p^2+M_B^2}(e^{\beta\sqrt{p^2+M_B^2}}-1)$, where $M_B^2=M^2+(2m+1)qB$, and $m$ denotes the Landau level.

In order to calculate the pion decay constant, we need to evaluate the matrix elements $\langle 0|\mathcal{A}_\mu^\pm|\pi^0\rangle$ and $\langle 0|\mathcal{A}_\mu^\pm|\pi^\mp\rangle$. At zero magnetic field, these are identical, but there are two pion decay constants at finite temperature; one for the time component and one for the spatial component of $\mathcal{A}_\mu$. The difference between them is an order-$p^4$ effect [37]. The matrix elements are proportional to $ip_\mu$ and the prefactors are denoted by $F_{\pi^0}$ and $F_{\pi^\pm}$, respectively. The expressions are divergent and requires renormalization of the $l_4$ and the renormalized result is

$$\begin{aligned}F_{\pi^0} &= F_\pi\left[1-\frac{1}{(4\pi)^2F^2}\left(I_H+J_1^B|qB|\right)\right] ,\\ F_{\pi^\pm} &= F_\pi\left[1-\frac{1}{2(4\pi)^2F^2}\left(I_H+J_1T^2+J_1^B|qB|\right)\right] ,\end{aligned}$$

where the pion decay constant $F_\pi$ in the vacuum is $F_\pi=(1+\frac{M^2}{(4\pi)^2F^2}\bar{l}_4)$. Note that these are different in a magnetic field since the contributions from the loops are different [8].

*Free energy and quark condensate:* We are interested in the contributions to the free energy that are due to a nonzero magnetic field and finite temperature. We therefore write the contribution to the free energy at the $n$'th loop order, $\mathcal{F}_n$, as a sum of three terms $\mathcal{F}_n=\mathcal{F}_n^{\text{vac}}+\mathcal{F}_n^B+\mathcal{F}_n^T$, where $\mathcal{F}_n^{\text{vac}}$ is the free energy in the vacuum, i. e. $B=T=0$, $\mathcal{F}_n^B$ is the zero-temperature contribution due to a finite magnetic field, and $\mathcal{F}_n^T$ is the finite-temperature contribution. The strategy is to isolate the term $\mathcal{F}_n^{\text{vac}}$ and subtract it from $\mathcal{F}_n$. This term contains ultraviolet divergences which are removed by renormalization of the low-energy constants of the chiral Lagrangian in the usual way and the renormalized $\mathcal{F}_n^{\text{vac}}$ represents vacuum energy of the theory. The sum $\mathcal{F}_n^B+\mathcal{F}_n^T$ generally contains ultraviolet divergences as well and it is rendered finite by renormalizing the $l_i$s. If we express the one-loop free energy in terms of $M_\pi$ instead of $M$, the dependence of $l_3$ cancels in the expressions for $\mathcal{F}^B_{1+2}$. and $\mathcal{F}^T_{1+2}$. This yields

$$
\begin{aligned}
\mathcal{F}^B_{1+2} &= \frac{M_\pi^4}{32\pi^2}\left[1 - 2\log\frac{M_\pi^2}{2|qB|}\right] + \frac{(qB)^2}{4\pi^2}\zeta^{(1,0)}(-1, \tfrac{1}{2} + x_\pi) + \frac{(qB)^2}{96\pi^2}\log\frac{\Lambda}{2qB} \\
&\quad -\frac{(qB)^2}{(4\pi)^4F^2}\left[\frac{1}{3}I_H(\bar{l}_6 - \bar{l}_5) + \bar{d}M^2\right] , \qquad (1) \\
\mathcal{F}^T_{1+2} &= -\frac{1}{32\pi^2}\left[J_0T^4 + 2J_0^B|qB|T^2\right] + \frac{M^2}{8(4\pi)^4F^2}\left[-J_1^2T^4 + 4J_1J_1^BT^2|qB| - 4I_HJ_1T^2\right] \\
&\quad -\frac{(qB)^3}{3(4\pi)^4F^2}(\bar{l}_6 - \bar{l}_5)J_1^B , \qquad (2)
\end{aligned}
$$

where $x_\pi = M_\pi^2/2|qB|$, $\bar{d}(M^2) = 8(4\pi)^4c_{34}^r + \frac{1}{3}(\bar{l}_6 - \bar{l}_5)\log\frac{\Lambda^2}{M^2}$, and $\Lambda$ is the renormalization scale. We note that the NLO correction to the free energy in the chiral limit does not vanish in contrast to the case of zero magnetic field.

At finite temperature, the quark condensate is $\langle\bar{q}q\rangle = \langle 0|\bar{q}q|0\rangle(1 + \frac{c}{F^2}\frac{\partial P}{\partial M_\pi^2})$, where the constant $c$ is defined by $c = -F^2\frac{\partial M_\pi^2}{\partial M}\langle 0|\bar{q}q|0\rangle^{-1}$ [7]. In the chiral limit, we have $c = 1$. In that case, the quark condensate reduces to

$$
\begin{aligned}
\langle\bar{q}q\rangle &= \langle 0|\bar{q}q|0\rangle\left\{1 + \frac{|qB|}{(4\pi)^2F^2}\ln 2 - \frac{|qB|}{(4\pi)^4F^2}\left[\frac{1}{3}(\bar{l}_6 - \bar{l}_5)(\log 2 + 1 - \gamma_E) - \bar{d}(qB)\right] - \frac{1}{2(4\pi)^2F^2}(J_1T^2 + 2J_1^B|qB|)\right. \\
&\quad \left. -\frac{T^2}{8(4\pi)^4F^4}\left(-J_1^2T^2 + 4J_1J_1^B|qB| - 4\ln 2\, J_1|qB|\right) - \frac{(qB)^3}{3(4\pi)^4F^2T^2}(\bar{l}_6 - \bar{l}_5)J_2^B\right\} , \qquad (3)
\end{aligned}
$$

where $M_B^2 = M^2 + (2m+1)qB$ and $J_2^B = -\beta^{-2}\sum_{m=0}^{\infty}\int_0^\infty p^{-2}dp/\sqrt{p^2 + M_B^2}(e^{\beta\sqrt{p^2+M_B^2}} - 1)$.

*Results and discussion:* We first notice that we in the limit $B \to 0$ recover the temperature dependence of $M_\pi$, $F_\pi$, $\mathcal{F}$, and $\langle\bar{q}q\rangle$ as in [5–7]. Similarly, we obtain the $T = 0$ result for the free energy and the $B$ dependence of the quark condensate as in [15–17]. We also notice that temperature dependence of the charged pion mass is the same as for vanishing magnetic field. The only difference is a temperature-independent constant proportional to $(qB)^2/F^2$.

In the remainder we focus on the chiral limit. It is then easy to show that $J_1T^2 \geq J_1^B|qB|$ for all values of $B$ and $T$. This implies that the pion decay constants $F_{\pi^0}$ and $F_{\pi^\pm}$ are larger than $F_\pi$. Moreover, for small values of $|qB|$, we can calculate the first corrections due to nonzero $B$ as a power series in $\sqrt{|qB|}/T$. One finds $F_{\pi^0} = F_\pi(1 - T^2/12F^2 + 5\sqrt{|qB|}T/48\pi F^2 + ...)$ and $F_{\pi^0} = F_{\pi^\pm}(1 - T^2/12F^2 + 5\sqrt{|qB|}T/96\pi F^2 + ...)$. Similarly, one can expand the quark condensate around $|qB| = 0$ and obtains $\langle\bar{q}q\rangle = \langle 0|\bar{q}q|0\rangle(1 + ...T^2/8F^2 + 5\sqrt{|qB|}T/96\pi F^2 + ...)$.

In Fig. 1, we show the quark condensate Eq. (3) as a function of temperature for $|qB| = 900$ (MeV)$^2$ (solid curve). For comparison, we also show the quark condensate for $|qB| = 0$ (dashed curve). We are using the experimental value $F_\pi = 93$ MeV. The quark condensate for vanishing $B$ goes faster to zero than it does in the presence of a magnetic field. This indicates that the critical temperature $T_c$ for the chiral transition is higher in a nonzero magnetic field. Of course, this conclusion is cautious since the behavior of the quark condensate in the vicinity of $T_c$ is beyond the reach of the low-temperature expansion. The result is in line with some model calculations, both mean-field type [29, 30, 33, 36] and beyond [34]. This is in disagreement with the recent lattice simulations [13] where a significant decrease of the crossover temperature was observed. On the other hand, the lattice result in Ref. [11] report a small increase of $T_c$ with $B$. Model calculations that seem to indicate a decrease can be found in Refs. [18, 32, 35]. A possible explanation for the different behavior of $T_c$ as a function of $B$ found in [32] and [33] could be the different treatment of the fermionic vacuum fluctuations and related renormalization issues.

In Ref. [18], the authors use ChPT to investigate the



FIG. 1: Quark condensate in the chiral limit as a function of temperature for $|qB| = 900$ (MeV)$^2$ (solid curve). and $|qB| = 0$ for comparison (dashed curve).

quark-hadron phase transition as a function of the magnetic field at the physical point. They compare the pressure in the hadronic phase with that of the quark-gluon plasma phase for an ideal gas of quarks and gluons, and subtracting the vacuum energy due to a nonzero gluon condensate $\langle g^2 G_{\mu\nu}G^{\mu\nu}\rangle$. For weak magnetic fields, the transition is first order. The line of first-order transitions ends at critical point. From this temperature onwards, the transition is a crossover. Normally, however, the critical temperature is determined by the behavior of the quark condensate. At the physical point, the condensate never vanishes and transition is a crossover. The crossover temperature is often defined by the inflection point of $\langle\bar{q}q\rangle$ as a function of temperature.

In conclusion, we have used chiral perturbation theory to calculate the pion masses, the decay constants, the free energy and the quark condensate at finite temperature in a magnetic background. Given the conflicting results for $T_c$ as a function of $B$ of various model calculations and lattice calculations, clearly more work needs to be done.

---

[1] S. Weinberg, Physica A **96**, 327 (1979).
[2] J. Gasser and H. Leutwyler, Ann. Phys. **158**, 142 (1984).
[3] J. Gasser and H. Leutwyler, Nucl. Phys. B **250**, 465 (1985).
[4] J. Bijnens, G. Colangelo, and G. Ecker, Ann. Phys. **280**, 100 (2000).
[5] J. Gasser and H. Leutwyler, Phys. Lett. B **184**, 83 (1987).
[6] J. Gasser and H. Leutwyler, Phys. Lett. B **188**, 477 (1987).
[7] P. Gerber and H. Leutwyler, Nucl. Phys. B **321**, 387 (1989).
[8] J. O. Andersen, in preparation.
[9] J. M. Lattimer and M. Prakash Phys. Rept. **442**, 109 (2007).
[10] D. E. Kharzeev, L. D. McLerran, and H. J. Warringa, Nucl. Phys. A **803** 227, 227 (2008).
[11] M. D'Elia, S. Mukherjee, and F. Sanfilippo, Phys. Rev. D **82**, 051501(R) (2010).
[12] M. D'Elia and F. Negro, Phys. Rev. D **83**, 114028 (2011).
[13] G. S. Bali, F. Bruckmann, G. Endrodi, Z. Fodor, S. D. Katz, S. Krieg, A. Schafer, and K. K. Szabo, e-Print: arXiv:1111.4956 [hep-lat].
[14] I. A. Shuspanov and A. V. Smilga, Phys. Lett. B **402**, 351 (1997).
[15] N. O. Agasian and I. A. Shushpanov, Phys. Lett. B **472**, 143 (2000).
[16] T. D. Cohen, D. A. McGady, and E. S. Werbos, Phys. Rev. C **76**, 055201 (2007).
[17] E. S. Werbos, Phys. Rev. C **77** (2008) 065202.
[18] N. O. Agasian and S. M. Fedorov, Phys. Lett. B **663**, 445 (2008).
[19] S. P. Klevansky and R. H. Lemmer, Phys. Rev. D **39**, 3478 (1989).
[20] V. P. Gusynin, V. A. Miransky, and I.A. Shovkovy, Phys. Lett. B **349**, (1995) 477; Nucl. Phys. B **462**, 249 (1996).
[21] A. Yu. Babansky, E. V. Gorbar, and G. V. Shchepanyuk, Phys. Lett. B **419**, 272 (1998).
[22] D. Ebert and K. G. Klimenko, Nucl. Phys. A **728**, 203 (2003).
[23] B. Hiller, A. A. Osipov, A. H. Blin, and J. da Providencia, SIGMA **4**, 024 (2008).
[24] J. K. Boomsma and D. Boer, Phys. Rev. D **81**, 074005 (2010).
[25] B. Chatterjee, H. Mishra, and A. Mishra, Phys. Rev. D **84**, 14016 (2011).
[26] S. S. Avancini, D. P. Menezes, and C. Providencia, Phys. Rev. C **83** 065805 (2011).
[27] M. Frasca and M. Ruggieri, Pys. Rev. D **83**, 094024 (2011).
[28] A. Rabhi and C. Providencia, Phys. Rev. C **83**, 055801 (2011).
[29] R. Gatto and M. Ruggieri, Phys. Rev. D **82**, 054027 (2010); ibid **83**, 034016 (2011).
[30] K. Kashiwa, Phys. Rev. D **83**, 117901 (2011).
[31] E. S. Fraga and A. J. Mizher, Phys. Rev. D **78**, 025016 (2008).
[32] J. O. Andersen and R. Khan, e-Print: arXiv:1105.1290 [hep-ph].
[33] A. J. Mizher, M. N. Chernodub, and E. S. Fraga, Phys. Rev. D **82**, 105016 (2010).
[34] V. Skokov, e-Print: arXiv:1112.5137 [hep-ph].
[35] E. S. Fraga and L. F. Palhares arXiv:1201.5881 [hep-ph]
[36] D. C. Duarte, R. L. S. Farias, and R. O. Ramos, Phys. Rev. D **84** (2011) 083525.
[37] R. D. Pisarski and M. Tytgat, Phys. Rev. D **540**, 2989 (1996).